

Secure Bidirectional Quantum Communication Protocol without Quantum Channel

Z. J. Zhang and Z. X. Man

Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China

*Email: zhangzj@wipm.ac.cn

(Dated: February 1, 2008)

In this letter we propose a theoretical deterministic secure direct bidirectional quantum communication protocol by using swapping quantum entanglement and local unitary operations, in which the quantum channel for photon transmission can be discarded, hence any attack with or without eavesdropping or even the destructive attack without scruple is impossible.

PACS Number(s): 03.67.Hk, 03.65.Ud

Much attention [1-6] has been focused on the study of the quantum key distribution (QKD) after the pioneering work of Bennett and Brassard published in 1984 [7], for the shared QKD can be used to encrypt the secret messages which is sent through a classical channel. As a matter of fact, the deterministic secure direct communication is more attractive and usually desired due to its obvious convenience. However, because of its more demanding on the security than QKDs, a proposal of a deterministic secure direct communication protocol is usually quite difficult. Till the end of 2003, only three deterministic secure direct communication protocols had been proposed by using the quantum entanglement of a photon pair [4-6]. Moreover, recently Zhang et al has proposed another deterministic secure direct communication protocol by using the quantum entanglement swapping of two photon pairs [8]. These four protocols mentioned are all message-unilaterally-transmitted protocols. Very recently, inspired by the deterministic secure direct protocol (i.e., the ping-pong protocol) proposed by Boström and Felbinger [5], a deterministic secure direct bidirectional simultaneous communication protocol is proposed by Zhang et al in a subtle way [9]. This is the first bidirectional secure quantum communication protocol. After this, according to the subtle idea presented in [9] the improvement on the two-step secure communication protocol [6] is also finished [10]. Hence, to our best knowledge, there are only six deterministic secure direct communication protocols so far. These protocols (except for the one in [8]) have four common properties as follows. (a) In all the protocols, after the message sender's encoding by unitary operation on the photon, the photon must be transmitted from the message sender's side to the message receiver's side via a quantum channel. This offers opportunities for the hostile person to eavesdrop or to attack the secret messages. (b) All the protocols are essentially quasisecure. Alternatively, the eavesdropping can not be detected with a possibility of 100%. As a result, a part of information might be leaked to the eavesdropper. (c) All the protocols are insecure under the attack without eavesdropping [11]. Hence, a strategy like message authentication should be adopted to detect such attacks. (d) In the cases that an eavesdropper is detected or an attacker attacks the quantum channel of photon transmission without scruple,

the secure communication has to be aborted. In fact, such cases occur quite possibly during some special times like a war. We think, all these are the common faults of the protocols, especially (d) is fatal. By the way, since a quantum channel for photon transmission should also be included in the protocol in [8], (d) is also the protocol's fault. In this letter, taking advantage of the idea of using the quantum entanglement swapping of two photon pairs [8] and the subtle idea presented in [9] to realize the bidirectional communication, we further propose another deterministic secure direct bidirectional simultaneous communication protocol, however, in which the quantum channel of photon transmission can be discarded, any attack with or without eavesdropping and even the destructive attack without scruple are accordingly impossible, hence this protocol is perfectly secure and can work in any case provided that the resource of the message carriers is sufficient. Consequently, this protocol will be very attractive in the commercial and military aspects due to its outstanding advantages.

Let us first describe the quantum entanglement swapping [12-14] simply. Let $|0\rangle$ and $|1\rangle$ be the horizontal and vertical polarization states of a photon, respectively. Then the four Bell states, $|\Psi^\pm\rangle = (|01\rangle \pm |10\rangle)/\sqrt{2}$ and $|\Phi^\pm\rangle = (|00\rangle \pm |11\rangle)/\sqrt{2}$, are maximally entangled states in the two-photon Hilbert space. Let the initial state of two photon pairs (i.e., the photon a_1 and b_1 pair and the photon a_2 and b_2 pair) be the product of any two of the four Bell states, such as $|\Psi_{a_1b_1}^+\rangle$ and $|\Psi_{a_2b_2}^+\rangle$, then after the Bell state measurements on the photon a_1 and a_2 pair and the photon b_1 and b_2 pair, since the following equation holds,

$$|\Psi_{a_1b_1}^+\rangle \otimes |\Psi_{a_2b_2}^+\rangle = \frac{1}{2}(|\Psi_{a_1a_2}^+\rangle|\Psi_{b_1b_2}^+\rangle - |\Psi_{a_1a_2}^-\rangle|\Psi_{b_1b_2}^-\rangle + |\Phi_{a_1a_2}^+\rangle|\Phi_{b_1b_2}^+\rangle - |\Phi_{a_1a_2}^-\rangle|\Phi_{b_1b_2}^-\rangle), \quad (1)$$

the total initial state (i.e., $|\Psi_{a_1b_1}^+\rangle \otimes |\Psi_{a_2b_2}^+\rangle$) is projected onto $|\eta_1\rangle = |\Phi_{a_1a_2}^+\rangle \otimes |\Phi_{b_1b_2}^+\rangle$, $|\eta_2\rangle = |\Phi_{a_1a_2}^-\rangle \otimes |\Phi_{b_1b_2}^-\rangle$, $|\eta_3\rangle = |\Psi_{a_1a_2}^+\rangle \otimes |\Psi_{b_1b_2}^+\rangle$ and $|\eta_4\rangle = |\Psi_{a_1a_2}^-\rangle \otimes |\Psi_{b_1b_2}^-\rangle$ with equal probability of $\frac{1}{4}$ for each. It is seen that previous entanglements between photons a_1 and b_1 , and a_2 and b_2 , are now swapped into the entanglements between photons a_1 and a_2 , and b_1 and b_2 . Therefore, if $|\Phi_{a_1a_2}^+\rangle$ is obtained by the Bell state measurements, $|\Phi_{b_1b_2}^+\rangle$ should be gained inevitably by the Bell state measurements; if $|\Phi_{a_1a_2}^-\rangle$ is obtained, then $|\Phi_{b_1b_2}^-\rangle$ is arrived at; and so on. This means that for a known initial state the Bell state measurement results after the quantum entanglement swapping are correlated. In the above example $|\Psi_{a_1b_1}^+\rangle \otimes |\Psi_{a_2b_2}^+\rangle$ is chosen as the initial state. In fact, similar results can also be arrived at provided that other choices of the initial states are given. As can be seen as follows:

$$\begin{cases} |\Psi_{a_1b_1}^+\rangle \otimes |\Psi_{a_2b_2}^-\rangle = \frac{1}{2}(|\Psi_{a_1a_2}^+\rangle|\Psi_{b_1b_2}^-\rangle - |\Psi_{a_1a_2}^-\rangle|\Psi_{b_1b_2}^+\rangle - |\Phi_{a_1a_2}^+\rangle|\Phi_{b_1b_2}^-\rangle + |\Phi_{a_1a_2}^-\rangle|\Phi_{b_1b_2}^+\rangle), \\ |\Psi_{a_1b_1}^+\rangle \otimes |\Phi_{a_2b_2}^+\rangle = \frac{1}{2}(|\Psi_{a_1a_2}^+\rangle|\Phi_{b_1b_2}^+\rangle - |\Psi_{a_1a_2}^-\rangle|\Phi_{b_1b_2}^-\rangle + |\Phi_{a_1a_2}^+\rangle|\Psi_{b_1b_2}^+\rangle - |\Phi_{a_1a_2}^-\rangle|\Psi_{b_1b_2}^-\rangle), \\ |\Psi_{a_1b_1}^+\rangle \otimes |\Phi_{a_2b_2}^-\rangle = \frac{1}{2}(|\Psi_{a_1a_2}^+\rangle|\Phi_{b_1b_2}^-\rangle - |\Psi_{a_1a_2}^-\rangle|\Phi_{b_1b_2}^+\rangle - |\Phi_{a_1a_2}^+\rangle|\Psi_{b_1b_2}^-\rangle + |\Phi_{a_1a_2}^-\rangle|\Psi_{b_1b_2}^+\rangle). \end{cases} \quad (2)$$

By the way, for the above four known initial states the correlation of the Bell state measurement results after the quantum entanglement swapping is very useful. On the other hand, it should also be noted that different results by the Bell state measurements correspond to different initial states for the above four known initial states. For examples, when $|\Psi_{a_1a_2}^+\rangle$ and $|\Psi_{b_1b_2}^-\rangle$ are obtained by the Bell state measurements, the initial state should be $|\Psi_{a_1b_1}^+\rangle \otimes |\Psi_{a_2b_2}^-\rangle$; when $|\Phi_{a_1a_2}^+\rangle$ and $|\Psi_{b_1b_2}^-\rangle$ are obtained by the Bell state measurements, the initial state should be $|\Psi_{a_1b_1}^+\rangle \otimes |\Phi_{a_2b_2}^-\rangle$; and so on. Incidentally, this property is used in our communication protocol. In addition, it

is easily verified that, the four Bell states can be transformed into each other by some unitary operations, which can be performed locally with nonlocal effects. For examples: Let $u_0 = |0\rangle\langle 0| + |1\rangle\langle 1|$, $u_1 = |1\rangle\langle 1| - |0\rangle\langle 0|$, $u_2 = |0\rangle\langle 1| + |1\rangle\langle 0|$, $u_3 = |0\rangle\langle 1| - |1\rangle\langle 0|$, then $|\Psi_{a_2b_2}^+\rangle$ will be in turn transformed into $|\Psi_{a_2b_2}^+\rangle$, $|\Psi_{a_2b_2}^-\rangle$, $|\Phi_{a_2b_2}^+\rangle$, $|\Phi_{a_2b_2}^-\rangle$ after the unitary operations u_0, u_1, u_2, u_3 on anyone photon (say, the b_2 photon) of the pair, respectively, that is, $u_0|\Psi_{a_2b_2}^+\rangle = |\Psi_{a_2b_2}^+\rangle$, $u_1|\Psi_{a_2b_2}^+\rangle = |\Psi_{a_2b_2}^-\rangle$, $u_2|\Psi_{a_2b_2}^+\rangle = |\Phi_{a_2b_2}^+\rangle$ and $u_3|\Psi_{a_2b_2}^+\rangle = |\Phi_{a_2b_2}^-\rangle$. Assume that each of the above four unitary operations corresponds to a two-bit encoding respectively, i.e., u_0 to '00', u_1 to '01', u_2 to '10' and u_3 to '11'. Then, taking advantage of the quantum entanglement swapping and the assumption of the two-bit codings, a deterministic secure direct bidirectional communication protocol can be proposed. We show it later.

Alice prepares an ordered N EPR photon pairs in state $|\Psi\rangle_{ab} = |\Psi^+\rangle = (|0\rangle_a|1\rangle_b + |1\rangle_a|0\rangle_b)/\sqrt{2}$ for each and divides them into two partner-photon sequences $[a_1, a_2, \dots, a_N]$ and $[b_1, b_2, \dots, b_N]$, where a_i (b_i) stands for the a (b) photon in the i th photon pair. Then Bob securely takes the b photon sequence away as a storage for the future use. By the way, as for how Bob can securely take the b photon sequence away as a storage, in principle, it is possible in theory. Maybe Bob can use some materials [15-16] to store the photons and take it away just as a baggage. Maybe the photons can be transmitted through a fiber to Bob's storage during the peaceful and secure times. In addition, how to maintain the entanglement properties of the photon pair is also a question. However, all these are only technological or other theoretical problems [17-18] and beyond our present theoretical scope. Whenever Alice and Bob want to communicate secretly with each other (if they want, they can publicly announce), they can do as follows. Both Alice and Bob perform the unitary operations on the photons with even (or odd) orders in their hands according to their secret messages. For examples, when Alice wants to let Bob securely know her bit string '011110...', according to her bit string she performs u_1, u_3, u_2 on the a_2, a_4 and a_6 photons respectively, and so on. When Bob wants to let Alice securely know his bit string '101100...', she performs u_2, u_3, u_0 on the b_2, b_4 and b_6 photons respectively, and so on. After their unitary operations they perform their Bell state measurements and publicly announces their results. That is, Alice performs in turn her Bell state measurements on the photon a_1 and a_2 pair, the photon a_3 and a_4 pair, etc, and publicly announce the measurement results in order; Similarly, Bob performs in turn his Bell state measurements on the photon b_1 and b_2 pair, the photon b_3 and b_4 pair, etc, and publicly announces the measurement results in order. Since Alice (Bob) knows which unitary operations she (he) has performed, then according to Bob's (Alice's) measurement results publicly announced and her (his) measurement results, she (he) can conclude Bob's (Alice's) unitary operations and accordingly extract Bob's (Alice's) bit string (See Table 1). So far a deterministic direct bidirectional communication has been proposed. By the way, if one party has no secret message to communicate, she or he can have two choices. One is that she or he only performs the unitary operations randomly to prevent from Eve's eavesdropping. The other choice is that, she (or he) publicly tells the partner that she (or he) has no secret message, then she (or he) does not perform any unitary operations and does not publicly announce the measurement results anymore. In any case of the above two choices, the communication protocol is reduced to a deterministic secure direct message-unilaterally-transmitted communication protocol.

Since Bob has taken securely the b photon sequence away and the a photon sequence is in Alice's hand, Eve (the eavesdropper) has no access to any photons, hence she can neither attack nor eavesdrop the secret messages. Therefore, the present deterministic direct bidirectional communication protocol is secure. In all protocols mentioned in this letter, the message carrier is photon. In all the other protocols except for [8], the photon encoded (i.e., performed a local unitary operation) has to be transmitted to the receiver to be performed a local Bell state measurement of the photon pair, hence the message transmission is essentially local. While in the present protocol, since the quantum entanglement swapping of two photon pairs is employed, the photons encoded need not to be transmitted via a quantum channel anymore, hence the message transmission is essentially nonlocal. It is this nonlocality which leads to the nonattackable property of the present protocol. This is the essential difference between the present protocol and other protocols. Incidentally, though in the present protocol the photon is chosen as the message carrier, the idea of the present protocol is a general one, hence it should also be suitable for other message carriers. Recently, the experimental achievement of Bose-Einstein condensates has attracted many attentions. It is reported that the entangled states of pairs of atoms can be created and the coherent transportation of the neutral atoms can be achieved in the optical lattices [19-20]. Maybe an alternative experimental demonstration of the present protocol by using the quantum entanglement swapping of the entangled atom pairs is feasible in the near future.

In summary, we have proposed a theoretical deterministic secure direct bidirectional quantum communication protocol by using swapping quantum entanglement and local unitary operations, in which the quantum channel for photon transmission can be discarded, hence any attack is impossible and accordingly the present protocol is perfectly secure.

This work is supported by the National Natural Science Foundation of China under Grant No. 10304022.

Table 1. Corresponding relations among Bob's and Alice's Bell state measurement results, the corresponding states before measurements and Bob's and Alice's

unitary u operations (i.e., the encoding bits).			
$\{\Phi_{a_1 a_2}^+, \Phi_{b_1 b_2}^+\}$	$\{\Psi_{a_1 a_2}^-, \Psi_{b_1 b_2}^+\}$	$\{\Psi_{a_1 a_2}^+, \Phi_{b_1 b_2}^+\}$	$\{\Psi_{a_1 a_2}^+, \Phi_{b_1 b_2}^-\}$
$\{\Psi_{a_1 a_2}^-, \Psi_{b_1 b_2}^-\}$	$\{\Phi_{a_1 a_2}^+, \Phi_{b_1 b_2}^-\}$	$\{\Psi_{a_1 a_2}^-, \Phi_{b_1 b_2}^-\}$	$\{\Psi_{a_1 a_2}^-, \Phi_{b_1 b_2}^+\}$
$\{\Psi_{a_1 a_2}^+, \Psi_{b_1 b_2}^+\}$	$\{\Phi_{a_1 a_2}^-, \Phi_{b_1 b_2}^+\}$	$\{\Phi_{a_1 a_2}^+, \Psi_{b_1 b_2}^+\}$	$\{\Phi_{a_1 a_2}^+, \Psi_{b_1 b_2}^-\}$
$\{\Phi_{a_1 a_2}^-, \Phi_{b_1 b_2}^-\}$	$\{\Psi_{a_1 a_2}^+, \Psi_{b_1 b_2}^-\}$	$\{\Phi_{a_1 a_2}^-, \Psi_{b_1 b_2}^-\}$	$\{\Phi_{a_1 a_2}^-, \Psi_{b_1 b_2}^+\}$
$\Psi_{a_1 b_1}^+ \otimes \Psi_{a_2 b_2}^+$	$\Psi_{a_1 b_1}^+ \otimes \Psi_{a_2 b_2}^-$	$\Psi_{a_1 b_1}^+ \otimes \Phi_{a_2 b_2}^+$	$\Psi_{a_1 b_1}^+ \otimes \Phi_{a_2 b_2}^-$
$\{u_0^A(00), u_0^B(00)\}$	$\{u_1^A(01), u_0^B(00)\}$	$\{u_2^A(10), u_0^B(00)\}$	$\{u_3^A(11), u_0^B(00)\}$
$\{u_1^A(00), u_1^B(00)\}$	$\{u_0^A(00), u_1^B(01)\}$	$\{u_0^A(00), u_2^B(10)\}$	$\{u_0^A(00), u_3^B(11)\}$
$\{u_2^A(00), u_2^B(00)\}$	$\{u_2^A(10), u_3^B(11)\}$	$\{u_1^A(01), u_3^B(11)\}$	$\{u_1^A(01), u_2^B(10)\}$
$\{u_3^A(00), u_3^B(00)\}$	$\{u_3^A(11), u_2^B(10)\}$	$\{u_3^A(11), u_1^B(01)\}$	$\{u_2^A(10), u_1^B(01)\}$

- [1] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Rev. Mod. Phys. **74**, 145 (2002).
- [2] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
- [3] D. Bruß, Phys. Rev. Lett. **81**, 3018 (1998).

- [4] A. Beige, B. G. Englert, C. Kurtsiefer, and H. Weinfurter, *Acta Phys. Pol. A* **101**, 357 (2002).
- [5] K. Bostrom and T. Felbinger, *Phys. Rev. Lett.* **89**, 187902 (2002).
- [6] F. G. Deng, G. L. Long, and X. S. Liu, *Phys. Rev. A* **68**, 042317 (2003).
- [7] C. H. Bennett and G. Brassard, in *Proceedings of the IEEE International Conference on Computers, Systems and Signal Processings, Bangalore, India* (IEEE, New York, 1984), p175.
- [8] Z. J. Zhang et al, qaunt-ph/0403218.
- [9] Z. J. Zhang et al, qaunt-ph/0403186.
- [10] Z. J. Zhang et al, qaunt-ph/0403215.
- [11] Q. Y. Cai, *Phys. Rev. Lett.* **91**, 109801 (2003).
- [12] J. W. Pan, M. Daniell, S. Gasparoni, G. Weihs, and A. Zeilinger, *Phys. Rev. Lett.* **86**, 4435 (2001).
- [13] M. Zukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, *Phys. Rev. Lett.* **71**, 4287 (1993).
- [14] S. Bose, V. Vedral, and P. L. Knight, *Phys. Rev. A* **57**, 822(1998).
- [15] D. F. Phillips, A. Fleischhauer, A. Maier, A. L. Walsworth, and M. D. Lukin, *Phys. Rev. Lett.* **86**, 783 (2001).
- [16] C. Liu, Z. Dutton, C. H. Behroozi, and L. V. Hau, *Nature (London)* **409**, 490 (2001).
- [17] H. J. Briegel, W. Dur, J. I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **81**, 5932 (1998).
- [18] Z. Zhao, T. Yang, Y. A. Chen, A. N. Zhang, and J. W. Pan, *Phys. Rev. Lett.* **90**, 207901 (2003).
- [19] L. M. Duan, A. Sorensen, J. I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **85**, 3991 (2000).
- [20] O. Mandel, M. Greiner, A. Widera, T. Rom, T. W. Hansch, and I. Bloch, *Phys. Rev. Lett.* **91**, 010407 (2003).